

L3 ANSWER 1 OF 4 MEDLINE
 AN 2001443764 IN-PROCESS
 DN 21382338 PubMed ID: 11489313
 TI Drug-polyionic block copolymer interactions for **micelle**
 formation: physicochemical characterisation.
 AU Govender T; Stolnik S; Xiong C; Zhang S; Illum L; Davis S S
 CS School of Pharmacy and Pharmacology, University of Durban-Westville,
 Private Bag, X54001, Durban, South Africa.
 SO JOURNAL OF CONTROLLED RELEASE, (2001 Aug 10) 75 (3) 249-58.
 Journal code: C46; 8607908. ISSN: 0168-3659.
 CY Netherlands
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS IN-PROCESS; NONINDEXED; Priority Journals
 ED Entered STN: 20010813
 Last Updated on STN: 20010813
 AB While covalent attachment of small drug molecules to AB copolymers for
 the formation of polymeric **micelles** for drug delivery has been
 investigated, few studies have focused on non-covalent interactions. The
 aim of this study was therefore to explore the potential of non-covalent
 interactions between an AB copolymer, Poly(aspartic acid)-poly(ethylene
 glycol) (Pasp-PEG), with anionic pendant groups and diminazene aceturate,
 a small molecular weight cationic drug. **Micelles** were prepared
 by mixing solutions of Pasp-PEG and diminazene in 25 mM Tris-HCl buffer.
 At all Pasp-PEG concentrations studied, the **micelles** appeared to
 be water soluble with a unimodal **size distribution** and
 ranged in size from approximately 22 to 60 nm. The polyionic
micelles also displayed similar and small absolute zeta potential
 values at various drug:monomer molar ratios which confirmed stabilisation
 by the PEG corona. The scattering intensity was maximal and remained
 unchanged, while particle size increased slightly at pH range from 3.4 to
 7.2. At this pH range both the polymer and drug would be ionised and
 ionic interactions possible to drive micellar formation. An increase in size
 and scattering intensity with addition of NaCl to the **micelles** was
 attributed to dehydration of the PEG corona which may have led to
 aggregation of the **micelles**. The absence of micellar
 dissociation upon addition of **salt** was attributed to the
 dominance of hydrogen bonding between Pasp and diminazene aceturate, as
 assessed by isothermal titration microcalorimetry. Morphological
 evaluation of these constructs showed them to be discrete and fairly
 uniform in size and shape. This study was therefore successful in
 confirming the potential of non-covalent interactions using an AB
 copolymer to form polyionic **micelles** for drug delivery.